27^e Congrès National

et

8^e Congrès Francophone
de réanimation

srlf

SOCIÉTÉ
DE RÉANIMATION
DE LANGUE FRANÇAISE

FAUT-IL TRAITER TOUTES LES HYPONATRÉMIES EN RÉANIMATION ?

Pr Ag Oussama JAOUED

Réanimation Médicale

CHU Taher Sfar Mahdia

Faut-il traiter toutes les hyponatrémies en réanimation ?

- Natrémie < 135mEq/L
- Trouble hydroélectrolytique le plus fréquent
- 20% des patients âgés de plus de 65 ans, 35% des hospitalisés, 30% des insuffisants cardiaques et 50% des patients avec une cirrhose ou cancer
- En réanimation, l'incidence des hyponatrémies varie de 10 à 45 % selon les études

2.1.1 Definition of hyponatraemia based on biochemical severity

We define 'mild' hyponatraemia as a biochemical finding of a serum sodium concentration ≤ 135 mmol/L

We define 'moderate' hyponatraemia as a biochemical finding of a serum sodium concentration ≤ 129 mmol/L

We define 'profound' hyponatraemia as a biochemical finding of a serum sodium concentration ≤ 125 mmol/L as measured by ion-selective electrode

2.1.2 Definition of hyponatraemia based on time of development

We define 'acute' hyponatraemia as hyponatraemia that is documented to exist for ≤ 48 h. We define 'chronic' hyponatraemia as hyponatraemia that is documented to exist for > 48 h. If there is no anamnestic evidence of acute hyponatraemia, we consider it being chronic

Table 1 (Table 5 of the online document): classification of symptoms of hyponatraemia

Severity	Symptom
Moderately severe	Nausea without vomiting Confusion Headache
Severe	Vomiting Cardio-respiratory distress Abnormal and deep somnolence Seizures Coma (Glasgow Coma Scale ≤ 8)

The epidemiology of intensive care unit-acquired hyponatraemia and hypernatraemia in medical-surgical intensive care units

Henry Thomas Stelfox^{1,2,3}, Sofia B Ahmed^{3,4}, Farah Khandwala⁵, David Zygun^{1,2,6}, Reza Shahpori¹ and Kevin Laupland^{2,1}

Table 3

Outcomes of care

Measures	Serum sodium category		
	Acquire hyponatraemia (n = 917)	Always normal (n = 5068)	p value [‡]
ICU length of stay, median (IQR), d [†]	6 (3 to 12)	2 (1 to 4)	<0.001
Hospital length of stay, median (IQR), d [†]	25 (14 to 50)	12 (7 to 24)	<0.001
ICU mortality, number (%)	164 (18)	456 (9)	<0.001
Hospital mortality, number (%)	255 (28)	799 (16)	<0.001

[†]ICU and hospital length of stay for patients who survive to hospital discharge.

[‡]p values calculated by multivariable linear and logistic regression for comparisons of patients who acquire hyponatraemia patients who always have normal serum sodium levels.

ICU = intensive care unit, IQR = interquartile range.

Fluctuations in Serum Sodium Level Are Associated With an Increased Risk of Death in Surgical ICU Patients*

- Prévalence à l'admission : 11.1% (n = 1,215/9431)
- Prévalence de l'hyponatrémie acquise en réanimation : 13.6% (n = 1,483)

TABLE 2. Characteristics and Outcome of the Study Cohort According to the Serum Sodium Level at Admission to the ICU

	Normonatremia n = 9431	Hyponatremia n = 1215
Mortality rates, n (%)		
ICU mortality	379 (4.0)	108 (8.9) ^d
Hospital mortality	701 (7.4)	201 (16.5) ^d

Figure 2. Bar chart representing ICU and hospital mortality rates according to sodium levels at admission to the ICU. $p < 0.01$ for ICU and hospital mortality among groups (Cochran–Armitage trend test).

Fluctuations in Serum Sodium Level Are Associated With an Increased Risk of Death in Surgical ICU Patients*

TABLE 3. Characteristics and Outcome of Patients With ICU-Acquired Dysnatremia and Those Who Remained Normonatremic During the ICU

	Normonatremia <i>n</i> = 6483	Hyponatremia <i>n</i> = 1483
Outcome, <i>n</i> (%)		
ICU mortality	91 (1.4)	48 (3.2) ^d
Hospital mortality	299 (4.6)	100 (6.7) ^c

Georg-Christian Funk
Gregor Lindner
Wilfred Druml
Barbara Metnitz
Christoph Schwarz
Peter Bauer
Philipp G. H. Metnitz

Incidence and prognosis of dysnatremias

Table 1 Unadjusted associa

Characteristic

Patients affected (%)
 Age (years), mean ± SD
 Males (%)
 Medical type of admission
 Scheduled surgical type of
 Unscheduled surgical type of
 Number of organ failures, n
 Sodium-corrected SAPS II s
 Observed ICU mortality (%)
 Length of ICU stay, days, n
 Observed hospital mortality

Fig. 2 Unadjusted (*open circles*) and adjusted (*filled circles*) risk of hospital mortality stratified by serum sodium on ICU admission. CI, confidence interval

stics

odium

(170)

P

<0.0001

<0.0001

<0.0001

<0.0001

<0.0001

<0.0001

<0.0001

<0.0001

<0.0001

<0.0001

Hyponatremia in critically ill patients

Rajesh Padhi, Baikuntha Nath Panda, Snehalata J...

- Etude prospective observationnelle
- 12 mois

- Groupe hyponatrémie (n=251) : 36%
- Hyponatremie à l'admission : 34%

- Hyponatrémie avec euvolémie : 148 (58.96%)
- Hyponatrémie avec hypervolémie : 66 (26.29%)
- Hyponatrémie avec hypovolémie : 37 (14.74%)

Table 2: Etiologic factors associated with hyponatremia

Disorders	No. of patients (n=251)	Percentage
SIADH* due to	91	36.25
Pneumonia	39	15.53
Drugs including SSRIs†, carbamazepine	15	5.97
Subarachnoid haemorrhage	5	1.99
After elective surgery	19	7.56
After emergency surgery	13	5.17
Severe sepsis	54	21.51
Trauma	53	21.11
Renal failure	15	5.97
Heart failure	13	5.17
Thiazides	12	4.78
Liver cirrhosis	9	3.58
Hypothyroidism	3	1.19
Hypocortisolism	1	0.37

Hyponatremia in critically ill patients

Rajesh Padhi, Baikuntha Nath Panda, Snehalata Jagati, Subhas Chandra Patra¹

Table 3: Outcomes and adverse events*

Outcome measure	Hyponatremia group (n=251)	Normal group (n=448)	Odds ratio or absolute difference (95% CI) [†]	Statistical test	P value
Death-no. of patients/total no. (%); all cause 30-day	49/251 (19.5)	74/448 (16.5)	0.4474 (0.3102-0.6454)	Mantel-Haenszel	0.0006
Days in hospital-median (IQR) [‡]	23 (15-23.5)	21 (14-23)	0	Log-rank test	0.15
Mechanical ventilation-no. of patients/total no. (%)	94/251 (37.45)	37/448 (8.25)	0.99 (0.62-1.58)	Pearson's test	<0.01
Days of mechanical ventilation	9.52±5.99	6.48±5.21	0	Wilcoxon rank-sum test	0.03
Renal-replacement therapy-no. of patients/total no. (%)	47/251 (18.7)	28/448 (6.3)	0.33 (0.12-0.87)	Pearson's test	0.03
Days of renal-replacement therapy	0.8±2.3	0.7±2.1	0	Wilcoxon rank-sum test	0.34

Moderate Hyponatremia Is A Risk of Mortality: Evidence from a Meta-Analysis

850222 patients dont 147948 patients avec hyponatrémie
 → 17,4% d'hyponatrémie

130.1±5.6 vs 134.9±5.1 mmol/L

Hyponatrémie était associée à la mortalité globale : RR = 2.60[2.31–2.93]; p= 0.0001

F

A**Odds ratio for overall mortality in myocardial infarction**

Source

Flear et al., 1979
 Goldberg et al., 20
 Goldberg et al., 2
 Klopotoski et al
 Havranek et al., 2
 Tada et al., 2011
 Tang et al., 2011

Overall**B**

Source

Panciroli et al., 198
 Adewole et al., 19
 Mien-Cheng et al.,
 VillaCorta et al., 2
 Gheorghide et al.,
 Gheorghide et al.,
 Milo-Cotter et al., 2
 Rusinaru et al., 200
 Barsheshet et al., 2
 DeWolfe et al., 20
 Baldasseroni et al.,
 Balling et al., 2011
 Shorr et al., 2011

Overall**C**

Source

Arroyo et al., 1976 (43)
 Vila et al., 1999 (44)
 Borroni et al., 2000 (45)
 Porcel et al., 2002 (46)
 Hackworth et al., 2009 (47)
 Radha Krishna et al., 2009
 Terg et al., 2009 (50)
 Jenq et al., 2010 (51)

Overall**Odds ratio for overall mortality in cirrhosis****D****Odds ratio for overall mortality in pulmonary infections**

Source

Singhi et al., 1992 (52)
 El-Ebiary et al., 1997 (54)
 Nair et al., 2007 (56)
 Song et al., 2008 (57)
 Zilberberg et al., 2008 (12)

Overall

No hyponatremia | hyponatremia

E

Odds ratio for overall mortality in mixed diseases

ORIGINAL RESEARCH

INFLUENCE OF EARLY DYSNATREMIA CORRECTION ON SURVIVAL OF CRITICALLY ILL PATIENTS

Michael Darmon,^{*†} Matthias Pichon,^{*†} Carole Schwebel,^{‡§} Stéphane Ruckly,^{‡§}
Christophe Adrie,^{||} Hakim Haouache,^{||} Elie Azoulay,^{**} Lila Bouadma,^{††}
Christophe Clec'h,^{‡‡} Maité Garrouste-Orgeas,^{§§} Bertrand Souweine,^{||||}
Dany Goldgran-Toledano,^{||||} Hatem Khallel,^{***} Laurent Argaud,^{†††‡‡‡}
Anne-Sylvie Dumenil,^{§§§} Samir Jamali^{||||||}, Bernard Allaouchiche,^{****}
Fabrice Zeni,^{*†} and Jean-François Thiebaud^{‡§}

SOCIÉTÉ
DE RÉANIMATION
DE LANGUE FRANÇAISE

Etude observationnelle

18 réanimations

Impact de la dysnatrémie sur la mortalité à j 28

A j01, 1,830 patients (25.9%) avaient une hyponatrémie :

- 1,187 patients (16.8%) d'hyponatrémie minime
- 435 patients (6.2%) d'hyponatrémie modérée
- 208 patients (2.9%) d'hyponatrémie sévère

1830 Patients with hyponatraemia at ICU admission
Hospital mortality: 342 (18.7%)

1260 Patients with hyponatraemia on day 3

811 Patients with persistent hyponatraemia
Hospital mortality: 163 (20.1%)

449 Patients with ICU-acquired hyponatraemia
Hospital mortality: 71 (15.8%)

INFLUENCE OF EARLY DYSNATREMIA CORRECTION ON SURVIVAL OF CRITICALLY ILL PATIENTS

Michael Darmon,^{*†} Matthias Pichon,^{*†} Carole Schwebel,^{‡§} Stéphane Ruckly,^{‡§}
 Christophe Adrie,^{||} Hakim Haouache,^{||} Elie Azoulay,^{**} Lila Bouadma,^{††}
 Christophe Clec'h,^{‡‡} Maité Garrouste-Orgeas,^{§§} Bertrand Souweine,^{||||}
 Dany Goldgran-Toledano,^{||||} Hatem Khallel,^{***} Laurent Argaud,^{†††‡‡‡}
 Anne-Sylvie Dumenil,^{§§§} Samir Jamali^{|||||}, Bernard Allaouchiche,^{****}
 Fabrice Zeni,^{*†} and Jean-François Timsit^{‡§}

SOCIÉTÉ
 DE RÉANIMATION
 DE LANGUE FRANÇAISE

TABLE 2. Factors independently associated with day 28 mortality

	OR	95% CI	P
Serum sodium concentration changes between day 1 and day 3			
Persistent hyponatremia	1.30	1.06 – 1.60	0.01

TABLE 3. Influence of serum sodium concentration changes on day 28 mortality after adjustment in patients with mild-to-severe dysnatremia

	OR	95% CI	P
Sodium correction rate, per mmol/L per day	0.93	0.90 – 0.97	0.0003

Hyponatremia Improvement Is Associated with a Reduced Risk of Mortality: Evidence from a Meta-Analysis

Giovanni Corona¹✉, Corinna Giuliani²✉, Joseph G. Verbalis³, Gianni Forti², Mario Maggi⁴, Alessandro Peri^{2*}

SOCIÉTÉ
DE RÉANIMATION
DE LANGUE FRANÇAISE

Studies included in quantitative synthesis
(meta-analysis)
N= 15

Hyponatremia Improvement Is Associated with a Reduced Risk of Mortality: Evidence from a Meta-Analysis

Giovanni Corona¹*, Alessandro Peri^{2*}

Fig 3. Odds ratio for overall mortality rate in patients with any increase of serum $[Na^+]$.

Hyponatremia Improvement Is Associated with a Reduced Risk of Mortality: Evidence from a Meta-Analysis

Giovanni C
Alessandro

- 08 études avec un seuil de correction > 130 mmol/l

Hyponatremia Improvement Is Associated with a Reduced Risk of Mortality: Evidence from a Meta-Analysis

Giovanni Corona¹✉, Corinna Giuliani²✉, Joseph G. Verbalis³, Gianni Forti², Mario Maggi⁴, Alessandro Peri^{2*}

- L'effet de l'amélioration de la natrémie sur la mortalité était plus évident :
 - Sujets âgés
 - Patient avec une natrémie plus basse à l'inclusion

**Faut-il traiter toutes les hyponatrémies
en réanimation ?**

OUI

COMMENT?

Diagnosis, Evaluation, and Treatment of Hyponatremia: Expert Panel Recommendations

Joseph G. Verbalis, MD,^a Steven R. Goldsmith, MD,^b Arthur Greenberg, MD,^c Cynthia Korzelius, MD,^d
Robert W. Schrier, MD,^e Richard H. Sterns, MD,^f Christopher J. Thompson, MD, FRCPI^g

^aGeorgetown University Medical Center, Washington, DC; ^bUniversity of Minnesota, Minneapolis, MN; ^cDuke University Medical Center, Durham, NC; ^dTufts University School of Medicine, Boston, MA; ^eUniversity of Colorado, Denver, CO; ^fUniversity of Rochester, Rochester, NY; ^gRoyal College of Surgeons in Ireland School of Medicine, Dublin, Ireland.

Intensive Care Med (2014) 40:320–331
DOI 10.1007/s00134-014-3210-2

GUIDELINES

Clinical practice guideline on diagnosis and treatment of hyponatraemia

Goce Spasovski, Raymond Vanholder¹, Bruno Allolio², Djillali Annane³, Steve Ball⁴,
Daniel Bichet⁵, Guy Decaux⁶, Wiebke Fenske², Ewout J Hoorn⁷, Carole Ichai⁸,
Michael Joannidis⁹, Alain Soupart⁶, Robert Zietse⁷, Maria Haller¹⁰, Sabine van der
Veer¹¹, Wim Van Biesen¹ and Evi Nagler¹ on behalf of the Hyponatraemia Guideline
Development Group

Table 1. Guideline-Recommended Treatment of Severely Symptomatic Hyponatremia With Hypertonic Saline

	US guideline	European guideline
Patients requiring treatment with hypertonic saline	Regardless of duration: <ul style="list-style-type: none"> • Hyponatremia with severe manifestations (somnolence, seizures, cardiorespiratory distress) • Hyponatremia with moderately severe symptoms (vomiting, confusion) in patients at high risk of progressing to life-threatening complications 	Regardless of duration: <ul style="list-style-type: none"> • Hyponatremia with severe manifestations (somnolence, seizures, cardiorespiratory distress) • Hyponatremia with moderately severe symptoms (vomiting, confusion) in patients at high risk of progressing to life-threatening complications
Rate of increase in serum sodium level desired (goal for change in serum sodium level)	4-6 mEq/L within 1-2 h	5 mEq/L within 1-2 h
Recommended treatment with hypertonic saline (3% sodium chloride)	100-mL bolus via central or peripheral vein over 10 min up to 3 times as needed to attain desired serum sodium level	150-mL bolus via central or peripheral vein over 20 min, repeating this step twice until the desired serum sodium level is achieved
Recommended frequency of measuring serum sodium level	After each bolus and every 4-6 h over the first 24 h	After each bolus and every 6 h over the first 24 h
Recommended increase in serum sodium level that should not be exceeded (correction limit)	At low risk for osmotic demyelination: <ul style="list-style-type: none"> • 10 mEq/L within first 24 h^a • 18 mEq/L within first 48 h^a At high risk for osmotic demyelination: <ul style="list-style-type: none"> • 8 mEq/L during any 24-h period^b 	At low risk for osmotic demyelination: <ul style="list-style-type: none"> • 10 mEq/L within first 24 h^a • 18 mEq/L within first 48 h^a At high risk for osmotic demyelination: <ul style="list-style-type: none"> • 8 mEq/L during any 24-h period^b

N Engl J Med 2015; 372:55-65

N Engl J Med 2000; 342:1581-1589

Table 3 Factors That Place Patients at High Risk of Developing the Osmotic Demyelination Syndrome with Correction of Chronic Hyponatremia

High Risk of Osmotic Demyelination Syndrome

- Serum sodium concentration ≤ 105 mmol/L
- Hypokalemia*
- Alcoholism*
- Malnutrition*
- Advanced liver disease*

Table 1. Single-center series of patients with hyponatremia: more rapid correction is associated with ODS

Author, ref., year	Patients	Serum	Patients	Correction	Correction rate	Correction
--------------------	----------	-------	----------	------------	-----------------	------------

Sterns [37],
Brunner et al [35], 1990
Tanneau et al [40], 1994
Ellis [36], 1990
Vu et al. [39], 1990

Continuous Versus Bolus Infusion of Hypertonic Saline in the Treatment of Symptomatic Hyponatremia Caused by SIAD

Aoife Garrahy,¹ Rosemary Dineen,¹ Anne Marie Hannon,¹ Martin Cuesta,¹ William Tormey,² Mark Sherlock,¹ and Chris J. Thompson¹

Comparer l'efficacité et l'innocuité de l'administration en bolus par rapport à la perfusion continue du NaCl 3% pour traiter les patients ayant un SIADH avec des symptômes sévères

Na⁺ < 125 mmol/l

Signes neurologiques : GCS <15, confusion, coma, convulsions

Continuous Versus Bolus Infusion of Hypertonic Saline in the Treatment of Symptomatic Hyponatremia Caused by SIAD

Table 2. Biochemical and Clinical Outcomes in Patients Receiving 3% Hypertonic Bolus or Continuous Infusion

	Bolus	Infusion	
	n = 22	n = 28	<i>P</i>
Change pNa, mmol/L			
6 h	6 (2–11)	3 (1–4)	<0.0001
12 h	8 (4–11)	5 (3–9)	<0.0001
24 h	10 (6–13)	10 (6–12)	NS
Change GCS			
6 h	3 (1–6)	1 (–2 to 2)	<0.0001
12 h	3 (1–6)	2 (–2 to 3)	0.0001
24 h	3 (1–7)	3 (1–6)	NS
Volume of 3% saline administered per 24 h, mL	200 (100–600)	500 (160–840)	<0.0001
pNa increase 8–12 mmol per 24 h, n	17	21	0.45
pNa increase >12 mmol per 24 h, n	1	0	0.48
Need for dextrose and/or dDAVP, n	5	0	0.008
Mortality, n	0	4	0.12

Treatment of symptomatic hyponatremia with hypertonic saline: a real-life observational study

Irina Chifu¹, Amelie Gerstl², Björn Lengenfelder^{2,3}, Dominik Schmitt^{2,3}, Nils Nagler^{2,3}, Martin Fassnacht¹ and Dirk Weismann²

N = 62 patients (mean sodium level: 118 mEq/L)
Bolus group (n = 36)

150 mL administered once for moderate symptoms or 150 mL administered twice for severe symptoms (per European guideline)

Comparator group (n = 26)
Standard care
Moderate symptoms

17 patients in bolus group vs 16 patients in comparator group (*P* value not provided)

Severe symptoms

19 patients in bolus group vs 10 patients in comparator group (*P* value not provided)

Observed change in serum sodium level

At 24 h: 9 mEq/L in bolus group vs 6 mEq/L in comparator group (difference not statistically significant, *P* value not provided)

Underwent relowering therapy for serum sodium level

27.8% in bolus group vs 0% in comparator group (*P* value not provided)

Osmotic demyelination

0 cases

Risk of Overcorrection in Rapid Intermittent Bolus vs Slow Continuous Infusion Therapies of Hypertonic Saline for Patients With Symptomatic Hyponatremia

The SALSA Randomized Clinical Trial

PRIMARY OUTCOME

Overcorrection at any given period, defined as increase in serum sodium level by >12 mmol/L within 24 h or >18 mmol/L within 48 h

N = 178 (mean serum sodium level: 118.2 mEq/L)

Bolus group (n = 87)

2 mL/kg administered 1 time for moderately severe symptoms or 2 mL/kg administered twice for severe symptoms

(per European guideline)

Continuous group (n = 91)

0.5 mL/kg/h administered for severe symptoms or administered for severe symptoms (adjustable rate for both doses)

Table 1. Baseline Characteristics of Intention-to-Treat Population

Characteristics	Patients, No. (%)		P value
	Rapid intermittent bolus (n = 87)	Slow continuous infusion (n = 91)	
Hyponatremic symptom			
Moderate	66 (75.9)	68 (74.7)	.86
Severe	21 (24.1)	23 (25.3)	

RCT: Risk of Overcorrection in Rapid Intermittent Bolus vs Slow Continuous Infusion for Symptomatic Hyponatremia

POPULATION

80 Men, 98 Women

Adults aged >18 y with symptoms of hyponatremia and glucose-corrected serum sodium ≤ 125 mmol/L

Mean (SD) age: 73.1 (12.2) y

INTERVENTION

178 Patients randomized

87 Rapid intermittent bolus

Hypertonic saline, 3%, infusion of 2 mL/kg over 20 min every 6 h

91 Slow continuous

Hypertonic saline, 3%, continuous infusion of 0.5-1 mL/kg/h

SETTINGS / LOCATIONS

3 General hospitals in the Republic of Korea

PRIMARY OUTCOME

Overcorrection at any given period, defined as increase in serum sodium level by >12 mmol/L within 24 h or >18 mmol/L within 48 h

FINDINGS

Both rapid intermittent bolus and slow continuous infusion for treating symptomatic hyponatremia are effective and safe with no difference in overcorrection risk (17.2% vs 24.2%, respectively; absolute risk difference, -6.9% (95% CI, -18.8% to 4.9%); $P = .26$)

Risk of Overcorrection in Rapid Intermittent Bolus vs Slow Continuous Infusion Therapies of Hypertonic Saline for Patients With Symptomatic Hyponatremia

The SALSA Randomized Clinical Trial

Table 2. Outcomes and Progressions in Intention-to-Treat Population

Variable	Patients, No. (%)		Absolute difference (95% CI)	P value
	Rapid intermittent bolus (n = 87)	Slow continuous infusion (n = 91)		
Post hoc analysis (efficacy as incidence of target correction rate by the time frame)				
sNa 5-9 mmol/L				
Within 1 h	28 (32.2)	16 (17.6)	14.6 (2 to 27.2) ^a	.02
Within 6 h	50 (57.5)	54 (59.3)	-1.9 (-16.4 to 12.6) ^a	.80
Within 24 h	78 (89.7)	79 (86.8)	2.8 (-6.6 to 12.3) ^a	.56
Mortality				
During admission	4 (4.6)	1 (1.1)	NA	.20
At 30 d	3 (3.4)	1 (1.1)	NA	.36

Risk of Overcorrection in Rapid Intermittent Bolus vs Slow Continuous Infusion Therapies of Hypertonic Saline for Patients With Symptomatic Hyponatremia

The SALSA Randomized Clinical Trial

Table 2. Outcomes and Progressions in Intention-to-Treat Population

Variable	Patients, No. (%)		Absolute difference (95% CI)	P value
	Rapid intermittent bolus (n = 87)	Slow continuous infusion (n = 91)		
Safety				
Incid				
Incid				

CONCLUSIONS AND RELEVANCE This randomized clinical trial found that both RIB and SIC therapies of hypertonic saline for treating hyponatremia were effective and safe, with no difference in the overcorrection risk. However, RIB had a lower incidence of therapeutic relowering treatment and tended to have a better efficacy in achieving sNa within 1 hour than SCI. RIB could be suggested as the preferred treatment of symptomatic hyponatremia, which is consistent with the current consensus guidelines.

Evaluation of Desmopressin in Critically Ill Patients with Hyponatremia Requiring 3% Hypertonic Saline

THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES
VOLUME 361 NUMBER 6 JUNE 2021

Association of intranasal desmopressin therapy with overcorrection

of severe
single-c

Naruhiko J
Mitsunaga

- Résultats des études non concluantes
- Rétrospectives
- Nécessités d'études randomisées

i Uenishi, MD^b
(2021) 53–61

Outcomes in Severe Hyponatremia Treated With and Without Desmopressin

Thomas E. MacMillan, MD, MSc, FRCPC,^{a,b,c} Rodrigo B. Cavalcanti, MD, MSc, FRCPC^{a,b,c}

THE AMERICAN
JOURNAL OF
MEDICINE®

Expert Panel Recommendation: Managing Excessive Correction of Chronic Hyponatremia

- Starting serum $[Na^+] \geq 120$ mmol/L: Intervention probably unnecessary.
- Starting serum $[Na^+] < 120$ mmol/L:
 - Replace water losses or administer desmopressin after correction by 6-8 mmol/L during the first 24 hours of therapy;
 - Withhold the next dose of vaptan if the correction is > 8 mmol/L;
 - Consider therapeutic re-lowering of serum $[Na^+]$ if correction exceeds therapeutic limits;
 - Consider administration of high-dose glucocorticoids (eg, dexamethasone, 4 mg every 6 hours) for 24-48 hours following the excessive correction.
- Re-lowering serum $[Na^+]$:
 - Administer desmopressin to prevent further water losses: 2-4 μ g every 8 hours parenterally;
 - Replace water orally or as 5% dextrose in water intravenously: 3 mL/kg/h;
 - Recheck serum $[Na^+]$ hourly and continue therapy infusion until serum $[Na^+]$ is reduced to goal (Figure 3).

Figure 2. Clinical Approach to Hypotonic Hyponatremia

ACUTE OR SEVERE SYMPTOMS?

YES

URINE SODIUM CONCENTRATION

≤ 30 mmol/L

> 30 mmol/L

≤ 100 mOsm/kg

LOW EFFECTIVE ARTERIAL BLOOD VOLUME

DIURETICS OR KIDNEY DISEASE?

YES

NO

Consider

- Primary polydipsia
- Low solute intake
- Beer potomania

If ECF expanded consider

- Heart failure
- Liver cirrhosis
- Nephrotic syndrome

If ECF reduced consider

- Diarrhea and vomiting
- Third spacing
- Remote diuretics

Consider

- Diuretics
- Kidney disease
- All other causes

If ECF reduced consider

- Vomiting
- Primary adrenal insufficiency
- Renal salt wasting
- Cerebral salt wasting
- Occult diuretics

If ECF normal consider

- SIAD
- Secondary adrenal insufficiency
- (Hypothyroidism)
- Occult diuretics

Hypovolemic Hyponatremia

- Kidney sodium losses
 - Diuretics
 - Osmotic diuresis (glucose, urea, mannitol)
 - Salt-wasting nephropathy
 - Adrenal insufficiency
 - Bicarbonaturia (renal tubular acidosis, disequilibrium stage of vomiting)
 - Ketonuria
 - Cerebral salt wasting syndrome
- Non-kidney sodium losses
 - Vomiting
 - Diarrhea
 - Blood loss
 - Excessive sweating
 - Fluid sequestration in the third space
 - Pancreatitis
 - Bowel obstruction
 - Peritonitis
 - Burns
 - Massive trauma

grès National

et

s Francophone

animation

srlf

SOCIÉTÉ
DE RÉANIMATION
DE LANGUE FRANÇAISE

Hypervolemic Hyponatremia

- Heart failure
- Cirrhosis
- Nephrotic syndrome
- Kidney failure (acute or chronic)
- Pregnancy

Euvolemic Hyponatremia

- Thiazide diuretics (certain cases)
- Severe hypothyroidism
- Glucocorticoid deficiency
- Low solute intake
 - Tea and toast diet
 - Beer potomania
 - Dilute infant formula
- Syndrome of inappropriate antidiuresis
 - Malignancies
 - Pulmonary and mediastinal
 - Nasopharyngeal
 - Gastrointestinal tract
 - Genitourinary tract
 - Drugs
 - Stimulants of synthesis of vasopressin: cocaine, nicotine, tricyclic antidepressants

antineoplastic agents (methotrexate, ifosfamide, cisplatin, carboplatin, vincristine, vinblastine, interferon), narcotics

- Arginine vasopressin-like compounds and enhancers of its action: desmopressin, oxytocin, vasopressin, vasopressin analogs
- Vasopressin receptor antagonists: conivaptan, tolvaptan
- Vasopressin synthetase inhibitors
- Mixed or unknown effects: cyclophosphamide, chlorpropamide, angiotensin-converting enzyme inhibitors, clofibrate, clozapine, selective serotonin reuptake inhibitors, 3,4-methylenedioxymethamphetamine, nonsteroidal anti-inflammatory drugs
- Central nervous system disorders
 - Mass lesions: tumor, abscess, hematoma
 - Inflammatory conditions: encephalitis, meningitis
 - Demyelinating disease: Guillain-Barré syndrome
 - Cerebrovascular accident
 - Head trauma
 - Acute psychosis
 - Delirium tremens
- Pulmonary conditions
 - Infections: bacterial or viral pneumonia, tuberculosis, aspergillosis
 - Asthma
 - Cystic fibrosis
 - Acute respiratory failure

- Chronic obstructive pulmonary disease
- Positive-pressure ventilation
- Other
 - Exercise-associated
 - Pain
 - Severe nausea
 - General anesthesia
 - Postoperative state
 - HIV/AIDS
 - Gain-of-function variants in the arginine vasopressin receptor 2 gene

Excessive Water Intake

- Primary polydipsia
- Multiple tap water enemas
- Fresh water drowning (eg, drowning in a lake or river)
- Dilute infant formula
- Exercise-associated
- Sodium-free irrigant solutions (hysteroscopy, laparoscopy, transurethral resection of the prostate)

Table 5 General Recommendations for Employment of Fluid Restriction and Predictors of the Increased Likelihood of Failure of Fluid Restriction

General recommendations:

- Restrict all intake that is consumed by drinking, not just water.
- Aim for a fluid restriction that is 500 mL/d below the 24-hour urine volume.
- Do not restrict sodium or protein intake unless indicated.

Predictors of the likely failure of fluid restriction:

- High urine osmolality (>500 mOsm/kg H₂O).
- Sum of the urine Na⁺ and K⁺ concentrations exceeds the serum Na⁺ concentration.
- 24-hour urine volume <1500 mL/d.
- Increase in serum Na⁺ concentration <2 mmol/L/d in 24-48 hours on a fluid restriction of ≤ 1 L/d.

Treatment of euvolemic hyponatremia in the intensive care unit by urea

Guy Decaux^{1*}, Caroline Andres¹, Fabrice Gankam Kengne¹, Alain Soupart^{1,2}

- 50 patients groupe hyponatrémie modérée
- $\text{Na}^+ = 128 \pm 4 \text{ mEq/L}$
- SIADH

- 35 patients groupe hyponatrémie sévère
- $\text{Na}^+ = 111 \pm 3 \text{ mEq/L}$

Figure 1 Evolution of SNa and blood urea in 50 patients before and after urea therapy.

Hyponatremia in critical care patients: Frequency, outcome, characteristics, and treatment with the vasopressin V₂-receptor antagonist tolvaptan[☆]

Bruce Friedman MD, CNSP, FCCP, FCCM^{a,b,*}, Joshua Cirulli PharmD^{c,*}

8. Potential uses of tolvaptan in critical care settings

Tolvaptan may have significant utility in treating hyponatremia in patients in a critical care setting (as noted previously, IV conivaptan is the preferred option for hospitalized patients who cannot tolerate oral medications; its use, however, is limited to a total of 4 days). In patients

Tolvaptan can be used to reverse the hyponatremia associated with thiazide diuretics in the euvolemic patient, as

blood urea nitrogen or creatinine levels. Tolvaptan treatment can also be used to address hyponatremia associated with many disease states seen in critically ill patients that are direct causes of SIADH, including serious infections, pneumonias or other lung injuries, and pain. Finally, it

association with symptomatic hyponatremia—a serious problem in the ICU—may potentially be reversed by correction of the hyponatremic state, tolvaptan is not indicated for use in this patient population (ie, patients requiring urgent intervention to prevent or treat serious neurological symptoms), nor has it been proved to provide symptomatic benefits.

Faut-il traiter toutes les hyponatrémies en réanimation ?

- OUI
- Lourde morbi-mortalité
- Hyponatrémie symptomatique : SSH 3% (recommandations américaine ou européenne)
- Hyponatrémie asymptomatique : traitement étiologique, restriction hydrique, urée? Tolvaptan?